# What are the Chances? A Look at Bayesian Network Modeling as a Predictive Decision Support Tool

PRESENTED AT THE 11TH NATIONAL MONITORING CONFERENCE


MARCH 25-29, 2019

DENVER, CO

DOUGLAS MCLAUGHLIN, KIESER & ASSOCIATES, LLC




## Decision-making & the NWQMC Monitoring Framework



### Water Resources Decisions Often Involve...

#### ▶ Numeric Thresholds

- Often create two or three categories or "conditions"
  - ▶ acceptable/not acceptable
  - good/fair/poor




### Water Resources Decisions Often Involve...

#### ▶ Numeric Thresholds

- Often create two or three categories or "conditions"
  - ▶ acceptable/not acceptable
  - good/fair/poor

#### <u>And</u> ► Causal Relationships

- ▶ How to achieve desired EPT status?
  - ▶ e.g., goal is EPT > 14
- How to support conditions for coldwater species?
  - e.g.,goal is coldwater macroinvertebrate taxa >= 4
- ► How to minimize excessive algal growth?
  - ▶ e.g., goal is chl a < 20 ug/L</p>




### Wrestling With Uncertainty

How to measure it. Its Impact on decision-making. How to communicate it.

Confidence In

It's About the Decision





## Common Questions for Decision Makers

#### **Common Management Questions**

What's our goal, and are we meeting it?


If not, what action should we take?

How confident are we that the system will respond as we think?

Can stakeholders clearly see what actions are being recommended and why?

## What Is A Bayesian Network Model?

- Bayes Theorem
- Conceptual models/influence diagrams
- Nodes (parent, child) and links showing relationships
- Bins/states, bin boundaries can reflect useful categories, thresholds
- Conditional probability tables (CPT) describing likelihoods of an outcome state given combinations of parent nodes
- Output in terms of probabilities (i.e., "what are the chances" of observing an output state/bin?)



From McLaughlin & Reckhow (2017)

$$P(A|B) = \frac{P(B|A) P(A)}{P(B)}$$

Example: A = EPT taxa; B = QHEI

### "Like A Drunk Uses A Lightpost..."

- "Statistics should be used like a drunk uses a lightpost...more for support than illumination." (or something like that. Dominique DiToro, at a conference sometime in the late '80s)
- ► The application here: BNs, like other modeling procedures, should be supported by a range of other analyses, including constructing a solid conceptual model and simply "poring over the data."

**Table 2**Spearman rank correlation coefficient for BN model endpoints (bold values p < 0.05).


	EPT	CHL	QHEI	MinDO	TP	TN
CHL	0.34					
QHEI	0.48	0.09				
MinDO	0.48	-0.02	0.16			
TP	-0.39	-0.25	-0.10	-0.19		
TN	0.16	0.00	0.11	0.31	0.56	
TKN	-0.62	-0.36	-0.25	-0.29	0.81	0.36

From McLaughlin & Reckhow (2017)

## BN Coldwater Macroinvertebrate Model (Ohio)

- ▶ Percent forest cover, mean water temperature, drainage area as predictors of coldwater macroinvertebrate taxa richness
   ≥4 in two ecoregions in Ohio
- CPTs generated using field data (954 records, expectationmaximization learning algorithm)
- Selection of explanatory variables and bin boundaries supported by other information and analyses
- One of the objectives: Provide a screening tool for identifying coldwater aquatic life use defined as coldwater taxa richness ≥4.

#### Calibrated Model




WAP = Western Alleghany Plateau; GLAC = Glaciated More information in Miltner & McLaughlin (2018)

## BN Coldwater Macroinvertebrate Model (Ohio)

#### Calibrated Model

#### totforest 0 to 20 20 to 58 31.4 58 to 100 39.0 $46 \pm 30$ COLD ALU 19.5 1 to 3 40.2 58.2 4 or more 23.0 7.34 meancelt $3.83 \pm 4.8$ 7.76 5 to 20 2.20 30.0 20 to 22 5.03 22 to 32 35.1 $20.1 \pm 6.8$ PCTCOLD **Ecoregion** 30.7 15 to 74 10.6 $9.24 \pm 14$ DRAINAGE 0.1 to 0.7 10.7 0.7 to 1.5 77.5 $2.69 \pm 1.4$

#### Model Scenario



### Sensitivity Analysis

- Mutual Information used to compare the relative influence of predictor variables for binned variables
  - Indicates how much having knowledge of one variable reduces the uncertainty in the value of another (equals 0 for independent variables)

**Table 4**Sensitivity analysis results by ecoregion for forest area, mean temperature, and drainage area.

Predictor	Ecoregions				
	Combined	WAP	GLAC		
	Mutual information (%)	Mutual information (%)	Mutual information (%)		
Forest area Drainage area Mean temperature	9.5 4.4 4	2.4 6.8 8.8	8.8 2.3 3.1		

WAP = Western Alleghany Plateau; GLAC = Glaciated From Miltner & McLaughlin (2018)

### CWH Model Performance: Model-Predicted Bin Vs. Test Data

K-fold validation using 80/20 train/test

Avg **Correct** Prediction >> 103/191 >> **54%** 

Zero	1 - 3	4 or more	Total	Actual
44 (39-49)	23 (18-28)	4 (0-8)	71	Zero
24 (20-29)	40 (34-47)	12 (2-21)	76	1 – 3
3 (1-5)	22 (8-35)	19 (9-29)	44	4 or more
71	85	35	191	Total


Avg "Worst" Prediction >> 7/191 = 4%

Zero	1 - 3	4 or more	Total	Actual
44 (39-49)	23 (18-28)	4 (0-8)	71	Zero
24 (20-29)	40 (34-47)	12 (2-21)	76	1 – 3
3 (1-5)	22 (8-35)	19 (9-29)	44	4 or more
71	85	35	191	Total


Spherical payoff = 0.66 (1.0 is best)

## Two Scenarios – The Influence of Forest Cover

▶ Less forest...fewer cold taxa



More forest...more cold taxa



One take-away: The model says that on average, the chances of finding 4 or more cold taxa are 3.4% when forest cover is 0-20% and 38.9% when forest cover exceeds 58%.

## BN Models as a Decision Support Tool: Questions Revisited

Common Management Questions	Relevant BN Model Characteristics
What's our goal, and are we meeting it?	Numeric environmental response targets/thresholds represented as bin boundaries
If not, what action should we take?	Conceptual models, causal inference, management options, sensitivity analyses
How confident are we that the system will respond as we think?	Scenario probabilities, model performance measures, updateable in adaptive management approach
Can stakeholders clearly see what actions are being recommended and why?	Graphic display of model components, interactive model scenarios with a single mouse click, transparent response variable targets, probability-based output

#### <u>Acknowledgements</u>

Robert Miltner, Kenneth Reckhow, National Council for Air and Stream Improvement, Inc.

### Questions?

#### References:

Marcot, B., Steventon, J., Sutherland, G., McCann, R., 2006. Guidelines for developing and updating Bayesian belief networks applied to ecological modeling and conservation. Can. J. For. Res. 36, 3063–3074.

McLaughlin, D.B., K.H. Reckhow. 2017. A Bayesian Network Assessment of Macroinvertebrate Responses to Nutrients and Other Factors in Streams of the Eastern Corn Belt Plains, Ohio, USA. Ecological Modelling. 345: 21-29. DOI: 10.1016/j.ecolmodel.2016.12.004.

Miltner, R., D.B. McLaughlin. 2018. Management of headwaters based on macroinvertebrate assemblages and environmental attributes. Science of the Total Environment 650: 438–451.